



Weierli Pet Product



HEBEI WEIERLI ANIMAL PHARMACEUTICAL GROUP CO.,LTD

PET CATALOGUE

Endoparasite	Wormer Clear (ivermectin Tablet)
	Cat Dewormer tablet
	Wormer Rid tablet
	Liquid Wormer (pyrantel Pamoate) Suspensions
	Wormer Wipe tablet
	Wormer Plus tablet
Ectoparasite	Nitenpyram tablet
	Spinosad tablet
	Amitraz 20% solution
	Natural Herbal Flea Tick Spray
Antibiotics	Enrofloxacin 68mg 136mh
	Metoclopramide
	Amoxicillin 250mg 500mg
	Gentamicin 250
Analgesics	Carprofen
Nutritional Supplements	Vitamin Supplement for Dog (tablet/powder)
	Chewable Calcium (tablet/powder)
	Healthy Coat (tablet/powder)
	Bone Live (tablet/powder)
	Liver Care (tablet/powder)
	Pet Blood producer (tablet/powder)
	Probiotic-vita (Paste/powder)
	Digestive Support powder
	Turmeric chewable tablet
	Pet milk

Endoparasite

Endoparasite



Wormer Wipe

Composition:

Each chewable tablet contains: Praziquantel 50 mg Pyrantel Pamoate ... 144 mg Febantel ... 150 mg Excipients...q.s

Uses:

In dogs: Treatment of mixed infections by nematodes and cestodes of the following species

Nematodes:

Ascarids: *Toxocara canis*, *Toxascaris leonina* (adult and late immature forms).

Hookworms: *Uncinaria stenocephala*, *Ancylostoma caninum* (adults).

Whipworms: *Trichuris vulpis* (adults).

Cestodes:

Tapeworms: *Echinococcus* species, (*E. granulosus*, *E. multilocularis*), *Taenia* species, (*T. hydatigena*, *T. pisiformis*, *T. taeniformis*), *Dipylidium caninum* (adult and immature forms).

Dosage: For oral administration only.

The recommended dose rates are: 15 mg/kg bodyweight febantel, 14.4 mg/kg pyrantel peanut and 5 mg/kg praziquantel. - 1 Praziquantel Tablet per 10 kg bodyweight;

Dosage Guide:

Puppies and small dogs

1/2-2 kg bodyweight = 1/4 Tablet 2-5 kg bodyweight = 1/2 Tablet

5-10 kg bodyweight = 1 Tablet

Dosage Guide:

Medium dogs

10-15 kg bodyweight = 1 1/2 Tablet 15-20 kg bodyweight = 2 Tablet

20-25 kg bodyweight = 2 1/2 Tablet 25-30 kg bodyweight = 3 Table

-For routine treatment a single dose is recommended.

Puppies should be treated at 2 weeks of age and every 2 weeks until 12 weeks of age. Thereafter they should be treated at 3 month intervals. It is advisable to treat the bitch at the same time as the puppies.

-For the control of *Toxocara*, nursing bitches should be dosed 2 weeks after giving birth and every two weeks until weaning.

Use During Pregnancy and Lactation:

-Consult a veterinary surgeon before treating pregnant animals for roundworms.

-The product may be used during lactation.

-Do not exceed recommended dose when treating pregnant bitches.

Contraindications, warnings, etc.:

-Do not use simultaneously with piperazine compounds.

-User safety: In the interests of good hygiene, persons administering the tablets directly to the dog, or by adding them to the dog's food, should wash their hands afterwards.

Special precautions for use

Special precautions for use in animals Any part used tablet should be discarded.

To ensure administration of a correct dose, body weight should be determined as accurately as possible Special precautions to be taken by the person administering the veterinary medicinal product to animals

In case of accidental ingestion, seek medical advice and show the package leaflet to the physician.

In the interests of good hygiene, persons administering the tablets directly to the dog, or by adding them to the dog's food, should wash their hands afterward



Wormer Plus



Ingredient

Febantel.....450mg
Pyrantel pamoate.....432mg
Praziquantel.....150mg
Ivermectina.....0,18mg

Directions

Wormer Plus contains Praziquantel, Pyrantel and Febantel making it the most effective dog wormer available.

Wormer Plus are uncoated flavoured dog worming tablets used in the treatment of mixed worm infestations in adult dogs and puppies. endogard treats Nematodes, Ascarids: *Toxocara canis*, *Toxascaris leonina* (late immature forms and mature forms), Hookworms: *Uncinaria stenocephala*, *Ancylostoma caninum* (adults), Cestodes *Taenia* spp., *Dipylidium caninum*.

Dosage

For most parasites, Wormer Plus given at a dose rate of 30kg. For cases of giardiasis 1 dose is recommended every 24 hours, for 3 days. Repeat the treatment at 30 days.

Contraindications and precautions: Do not administer in pregnant females. Sometimes vomiting, diarrhea, depression and salivation may occur.

If you suspect that your pet has received an overdose of Wormer Plus , consult your veterinarian or emergency veterinary clinic immediately.

Environmental Precautions:

Any unused product or waste material should be disposed of in accordance with current national requirements.

Pharmaceutical Precautions:

No special storage precautions.

Operator Precautions:

None

General Precautions:

For animal treatment only
Keep out of reach of children



CATS DEWORMER

BROAD SPECTRUM DEWORMER FOR CATS AND KITTENS

Praziquantel/pyrantel pamoate Tablets

Ingredient:**Active substances:** Pyrantel pamoate 230 mg Praziquantel 20 mg

Indications for use, specifying the target species

For the treatment of mixed infections caused by the following gastrointestinal roundworms and tapeworms:

Roundworms: *Toxocara cati*, *Toxascaris leonina*.**Tapeworms:** *Dipylidium caninum*, *Taeniataeniaeformis*,**The recommended dose :** 20 mg/kg pyrantel (57.5 mg/kg pyrantel embonate) and 5 mg/kg praziquantel. This is equivalent to 1 tablet per 4 kg bodyweight.

Body weight	Tablets
1.0 - 2.0 kg	½
2.1 - 4.0 kg	1
4.1 - 6.0 kg	1 ½
6.1 - 8.0 kg	2

Caution :

KEEP OUT OF REACH OF CHILDREN. Consult your veterinarian before administering to sick or pregnant animals.

STORAGE CONDITIONS: Store at 20°C-25°C (68°F-77°F), excursions permitted between 15°C-30°C (59°F-86°F). Brief exposure to temperatures up to 40°C (77°F); however such exposure should be minimized.

LIQUID WORMER (PYRANTEL PAMOATE) SUSPENSIONS

Parasiticide-Oral For Puppies And Kitties

INDICATIONS:

Pyrantel Pamoate is used to treat parasites such as roundworms and hookworms in dogs and cats. Most puppies and kittens are born with internal parasites or worms derived from their mother. These parasites may injure the pet and pose health risks to children who might eat soil contaminated with the feces of infected dogs or cats. Accordingly, veterinarians and public health officials advise pet owners to deworm puppies and kittens during the first few months of life.

Pyrantel pamoate is one of the most commonly used drugs for de-worming puppies and kittens. It can also be used for parasite control in adult pets and is relatively safe when administered to sick or debilitated animals requiring de-worming.

Pyrantel pamoate acts on the nervous system of certain parasites, resulting in paralysis and death of the worm.

Pyrantel pamoate may also be used to prevent reinfestation of *Toxocara canis* in puppies and adult dogs and in lactating bitches after whelping.

NET WEIGHT: 45ml

ACTIVE INGREDIENT: 4.54mg of pyrantel base as pyrantel pamoate per ml.(Easy to administer shake well before use.

DIRECTIONS FOR USE:

Administer 1 full teaspoon (5 ml) for each 10 lb of body weight. To assure proper dosage, weigh animal prior to treatment. If there is reluctance to accept the dose, mix in a small quantity of dog food to encourage consumption.

It is recommended that dogs maintained under conditions of constant exposure to worm infestation should have a follow-up fecal exam within 2-4 weeks after first treatment.

If your dog looks or acts sick, consult your veterinarian before treatment. For maximum control and prevention of reinfestation, it is recommended that puppies be treated at 2, 3, 4, 6, 8 and 10 weeks of age. Lactating bitches should be treated 2-3 weeks after whelping. Adult dogs kept in heavily contaminated quarters may be treated at monthly intervals to prevent *Toxocara canis* reinfestation.

PRECAUTION AND SIDE EFFECTS

While generally safe and effective when prescribed by a veterinarian, pyrantel pamoate can cause side effects in some animals.



Endoparasite

- Pyrantel pamoate should not be used in animals with known hypersensitivity or allergy to the drug.
- Pyrantel pamoate is well tolerated by most sick animals and is one of the safest de-wormers for use in puppies and kittens. However, use should be avoided in severely ill animals if there is no indication for de-worming.
- If given at an appropriate dose, adverse effects are rare.
- A small percentage of animals vomit after receiving pyrantel pamoate.

RECOMMENDED STORAGE: Recommended Storage: Store Below 30°C (86°F)

ENVIRONMENTAL PRECAUTIONS:

Any unused product or waste material should be disposed of in accordance with current national requirements.

PHARMACEUTICAL PRECAUTIONS:

No special storage precautions.

OPERATER PRECAUTIONS:

None

GERENALPRECAUTIONS:

For animal treatment only
Keep out of reach of children

WORM CLEAR IVERMECTION TABLET

Endoparasite



Review of Ivermectin for Dogs and Cats

Ivermectin, also known as ivermectin, is used to control skin parasites, gastrointestinal parasites and parasites within the bloodstream in dogs and cats.

Parasitic diseases are common in animals. Parasites can affect the skin, ears, stomach and intestines, and the internal organs including the heart, lungs and liver. Several drugs have been developed to kill or prevent parasites such as fleas, ticks, mites and worms. Ivermectin and related drugs are among the most effective of these.

Ivermectin is a parasite control drug. Ivermectin causes neurologic damage to the parasite, resulting in paralysis and death.

Ivermectin has been used to prevent parasite infections, as with heartworm prevention, and to treat infections, as with ear mites.

Ivermectin is a prescription drug and can only be obtained from a veterinarian or by prescription from a veterinarian.

Dosing Information of Ivermectin for Dogs and Cats

Medication should never be administered without first consulting your veterinarian. The dose for ivermectin varies from species to species and also depends on the intent of treatment. General dosing guidelines follow.

For dogs: Dose is 0.0015 to 0.003 mg per pound (0.003 to 0.006 mg/kg) once a month for heartworm prevention; 0.15 mg per pound (0.3 mg/kg) once, then repeat in 14 days for skin parasites; and 0.1 mg per pound (0.2 mg/kg) once for gastrointestinal parasites.

For cats: Dose is 0.012 mg per pound (0.024 mg/kg) once monthly for heartworm prevention.

The duration of administration depends on the condition being treated, response to the medication and the development of any adverse effects. Be certain to complete the prescription unless specifically directed by your veterinarian. Even if your pet feels better, the entire treatment plan should be completed to prevent relapse or prevent the development of resistance.

WORM CLEAR IVERMECTION TABLET

Side Effects of Ivermectin in Cats :

In cats, ivermectin has a fairly high margin of safety. When seen, side effects include:

- Agitation
- Crying
- Lack of appetite
- Dilated pupils
- Paralysis of hind legs
- Muscle tremors
- Disorientation
- Blindness
- Other neurological signs, such as head pressing or wall climbing

If your cat is receiving ivermectin and you notice these types of symptoms, discontinue the medication and contact your veterinarian.

Side Effects of Ivermectin in Dogs :

In dogs, the risk of side effects associated with ivermectin depends on the dosage, on the susceptibility of the individual dog and on the presence of heartworm microfilaria (a larval form of the heartworm.)

When used at a low dose for heartworm prevention in a dog free of heartworms, ivermectin is relatively safe. At higher doses which may be used for treating other parasitic infections, the risk of side effects increases.

Potential side effects include:

- Vomiting
- Dilated pupils
- Muscle tremors
- Blindness
- In-coordination

- Lethargy
- Lack of appetite
- Dehydration

When used in a dog infected with heartworms, a shock-like reaction believed to be caused by dying microfilaria can occur. This type of reaction may be accompanied by lethargy, a low body temperature and vomiting. Dogs testing positive for heartworms should be observed closely for at least 8 hours following the administration of ivermectin.

Ivermectin Sensitivity in Collies and Similar Breeds :

Neurotoxicity can also occur with ivermectin usage in some dogs. This is particularly common in dogs that have a genetic mutation known as the MDR1 (multi-drug resistance) gene mutation. This gene mutation is known to occur most commonly in breeds such as Collies, Australian Shepherds, Shelties, Long-haired Whippets and other breeds with "white feet."

Notice:

- Ivermectin should not be used in animals with known hypersensitivity or allergy to the drug.
- Ivermectin should not be used in dogs that are positive for heartworm disease except under strict supervision of a veterinarian.
- Prior to starting a heartworm prevention containing ivermectin, the dog should be tested for heartworms.
- Ivermectin generally should be avoided in dogs less than 6 weeks of age.

Worm Rid

PRAZIQUANTEL FENBENDAZOLE TABLET

FOR DOGS & CATS



Product characteristics:

Fenbendazole is a drug prescribed by veterinarians to treat intestinal parasites. It kills roundworms, whipworms, hookworms, and tapeworms in pets. It is only FDA-approved in dogs, but veterinarians often prescribe Fenbendazole to cats as well. Fenbendazole works by interrupting the parasites' cellular transport and metabolism. They diminish the parasites' energy reserves and limit their ability to eliminate waste and produce protective factors from their cells.

Praziquantel is effective against liver flukes and tapeworms. Tapeworms are spread by ingesting a flea or louse that has in turn ingested a tapeworm egg. A flea preventative may also be prescribed in conjunction with administration of Praziquantel to prevent any future tapeworm infections.

Indication:

A broad spectrum anthelmintic for the treatment of mixed infections of gastrointestinal nematodes and cestodes in dogs and cats.

Roundworms Ascarids *Toxocara canis*, (immature, adult), *Toxocara cati*, (adult), *Toxascaris leonina* (immature, adult)

Hookworms *Uncinaria stenocephala*, (immature, adult), *Ancylostoma caninum* (immature, adult)

Whipworms *Trichuris vulpis* (adult)

Tapeworms *Echinococcus granulosus*, *Echinococcus multilocularis*, *Dipylidium caninum*, *Taenia* spp., *Mesocestoides* spp

This product may also be used as an aid in the control of *Giardia* protozoa in dogs and *Aelurostrongylus abstrusus* lungworm infection in cats.



Composition:

Each tablet contains

Praziquantel 50mg

Fenbendazole 500mg

Dosage and Administration:

This product is administered orally either directly or mixed with a portion of meat or sausage or mixed with food. Dietary measures or fasting are not necessary.

Absorption may be improved with food.

Worm Rid PRAZIQUANTEL FENBENDAZOLE TABLET

FOR DOGS & CATS

Routine Treatment of adult dogs:

This product should be administered as a single treatment at a dose rate of 5 mg praziquantel and 50 mg fenbendazole per kg bodyweight (equivalent to 1 tablet per 10 kg).

For example:

Small dogs and puppies over 6 months of age

0.5 - 2.5 kg bodyweight 1/4 tablet

2.5 - 5 kg bodyweight 1/2 tablet

6 - 10 kg bodyweight 1 tablet

Medium sized dogs

11 - 15 kg bodyweight 1 1/2 tablets

16 - 20 kg bodyweight 2 tablets

21 - 25 kg bodyweight 2 1/2 tablets

26 - 30 kg bodyweight 3 tablets

Large Dogs

31 - 35 kg bodyweight 3 1/2 tablets

36 - 40 kg bodyweight 4 tablets

Routine Treatment of adult cats:

This product should be administered as a single treatment at a dose rate of 5 mg praziquantel and 50 mg fenbendazole per kg bodyweight (equivalent to 1/2 tablet per 5 kg bodyweight)

For example:

0.5 - 2.5 kg bodyweight 1/4 tablet

2.5 - 5 kg bodyweight 1/2 tablet

For routine control adult dogs and cats should be treated once every 3 months.

Weaned puppies & kittens under 6 months of age:

This product should be administered at a dose rate of 5 mg praziquantel and 50 mg fenbendazole per kg bodyweight (equivalent to 1/2 tablet per 5 kg bodyweight). Treatment should be administered for three consecutive days.

Unweaned puppies and nursing bitches:

For the control of Toxocara, it is important to worm young puppies very regularly with the product, at a dose rate of 5 mg praziquantel and 50 mg fenbendazole per kg bodyweight daily for three consecutive days (equivalent to 1/2 tablet per 5 kg daily for 3 days). This treatment regimen should be repeated at 2 weekly intervals from the age of 2 weeks for pups less than 12 weeks of age.

It is then recommended that this product be administered at intervals of 3 months.

Nursing bitches should be treated at the same time and as frequently as puppies up to 12 weeks of age. Thereafter the adult worming regime of once every three months is recommended.

Increased dosing for specific infections:

For the treatment of Clinical worm infestations in adult dogs administer this product at a dose rate of:

5mg praziquantel and 50mg fenbendazole per kg bodyweight daily for two consecutive days (equivalent to 1 tablet per 10 kg daily for 2 days)

For the treatment of Clinical worm infestations in adult cats and as an aid in the control of the Lungworm, Aelurostrongylus abstrusus in cats and Giardia protozoa in dogs administer this product at a dose rate of:

5 mg praziquantel and 50 mg fenbendazole per kg bodyweight daily for three consecutive days (equivalent to 1/2 tablet per 5 kg daily for 3 days).

Note:

Not intended for use in kittens less than 8 weeks of age.

Do not exceed the stated dose when treating pregnant bitches.

A veterinary surgeon should be consulted before treating pregnant bitches for roundworm.

Do not use in pregnant cats.

Safe for use in lactating animals. Both fenbendazole and praziquantel are very well tolerated.

After severe overdose occasional vomiting and transient diarrhoea may occur. Inappetence may occur following high doses in cats.

Ectoparasite

Ectoparasite



Spinosad Oral Tablet for Pet

This is chewable beef-flavored tablet that kills fleas and prevents flea infestations on dogs and cats for a full month. for Dogs & Cats requires a prescription from your veterinarian.

For:

Dogs & Cats (14 weeks of age or older)

Benefits:

- Starts killing fleas within 30 minutes
- Lasts a full month
- Kills fleas before they can lay eggs
- Comes in a convenient chewable, beef-flavored tablet
- Spinosad, the active ingredient in , is environmentally friendly

How it Works:

Spinosad kills fleas by causing excitation of the insect's nervous system, leading to paralysis and rapid death.

Cautions:

Side effects may occur, including vomiting. If your pet is experiencing loss of appetite, lethargy, or any other difference in behavior, contact your veterinarian. Dogs receiving extra-label doses of ivermectin to treat non-responsive demodectic mange and other conditions have developed signs of ivermectin toxicity. Ivermectin is an ingredient found in heartworm preventatives such as Heartgard Plus, Heartgard, Iverhart Plus and Ivomec. In cats, the most common adverse reaction recorded in clinical trials was vomiting. Other adverse reactions were: lethargy, decreased appetite, weight loss, and diarrhea. Use with caution with concomitant extra-label use of ivermectin.

Generic Name:

Spinosad

How should this medication be given:

Give chewable tablets with food. Give chewable tablets once a month. chewable tablets may be chewed, offered in food, or administered like other tablet medications. If vomiting occurs within 1 hour of administration, redose the pet with another full dose. Store at room temperature

What are the possible side effects :

Side effects that may occur include vomiting. If vomiting occurs within one hour of administration, redose with another full dose. Other side effects may occur. Continue giving chewable tablets and talk to your veterinarian if your pet experiences decreased appetite, lethargy or decreased activity, diarrhea, cough, increased thirst, vocalization, increased appetite, redness of the skin, hyperactivity and excessive salivation. Talk to your veterinarian about any side effect that seems unusual or especially bothersome to the pet.

What other drugs will affect :

Talk to your veterinarian before giving any prescription or over the counter medications.

Ectoparasite

Nitenpyram Oral Tablet for Pet

Indication

Nitenpyram Tablets kill adult fleas and are indicated for the treatment of flea infestations on dogs, puppies, cats and kittens 4 weeks of age and older and 2 pounds of body weight or greater. A single dose of Nitenpyram should kill the adult fleas on your pet. If your pet gets re-infested with fleas, you can safely give another dose as often as once per day.

Dosage

Nitenpyram is an oral tablet for dogs, puppies, cats and kittens 4 weeks of age and older and 2 pounds of body weight or greater.

Formula	Pet	Weight	Dose
11.4 mg	Dog or Cat	2-25 lbs	1 Tablet
57.0 mg	Dog	25.1-125 lbs	1 Tablet

Active Ingredients

Nitenpyram 11.4 mg

Warnings

Not for human use. Keep this and all drugs out of the reach of children.

Directions

To give Nitenpyram Tablets, place the pill directly in your pet's mouth or hide it in food. If you hide the pill in food, watch closely to make sure your pet swallows the pill. If you are not sure that your pet swallowed the pill, it is safe to give a second pill. Treat all infested pets in the household. Fleas can reproduce on untreated pets and allow infestations to persist.

Storage Conditions

Store at controlled room temperature, between 59 °F and 77 °F (15°C-25°C).



Amitraz 20% solution Dogs

Long Lasting Ectoparasite Control

Product Description

It is a pale yellow clear liquid. This is an insecticide and expellant, used for control of ectoparasites, such as mites, ticks and lice etc. for goat, sheep, camel, cattle, pigs, and it is also suitable for dogs, cats and rabbits. It is relative safe to predatory insects and bees. No drug accumulating in the tissues of animal. Amitraz 20% solution is specially formulated for animal ectoparasite control. Its exceptional activity against a broad spectrum of ticks, mange mites, lice and keds has been demonstrated in field trials and commercial use throughout the world. Three major indications exist; cattle tick control, pig mange and sheep mange.

**KEEP OUT OF THE REACH OF CHILDREN
FOR VETERINARY USE ONLY**

Administration and dosage:

External use only.

Animal specials	Application	dilution	replenishment	interval
Dogs	Ticks & mange	10ml : 4L water	1L : 2L water	10 days

Control of all stages of tetranychid and eriophyid mites, pear suckers, scale insects, mealybugs, whitefly, aphids, and eggs and first instar larvae of Lepidoptera on pome fruit, citrus fruit, cotton, stone fruit, bush fruit, cotton, stone fruit, bush fruit, strawberries, hops, cucurbits, aubergines, capsicums, tomatoes, ornamentals.

Withholding Period:

Taken into consideration the maximum residual limits the following withdrawal periods are recommended. For edible tissues; 1 day for cattle and goats and 7 days for sheep and pig. For bulked milk from treated herbs a zero withholding period can be recommended provided that least 7 hours have elapsed between treatment and the next milking.

USE DURING PREGNANCY, LACTATION OR LAY

Side effects

Apart from occurrences of dose related somnolence in cattle and sheep, which disappear spontaneously after a few hours, Amitraz 20% solution has an excellent safety margin and can be safely used in farm animals any stage of pregnancy.

Each 1000 Litres contains

Active ingredient(s) and amount(s): Each ml contains

Amitraz 200mg

Warning: This product should not be used for horse because it can cause irreversible gut stasis.

Storage: Keep jar tightly closed to preserve freshness at below 30 C

Package: 100ml

Natural Herbal Flea Tick Spray

Description

When fleas and ticks have your home and dog under assault, Vet's Best Dog Flea + Tick Home Spray fights back with its unique blend of certified natural essential oils and plant-based ingredients. Peppermint oil and clove extract (eugenol) help control and kill fleas, flea eggs, and ticks on contact, and also repels mosquitoes too! It uses absolutely no chemical pyrethrins, permethrins or cedar oil, so it's safe for the whole family. The non-staining formula is great for home surfaces like furniture, bedding, carpets, flooring and countertops, so it can be used everywhere you need it. Safe for use on or around dogs and puppies 12 weeks or older.

Ingredients

Peppermint Oil 0.30%, Eugenol (Clove Extract) 0.70%, Sodium Lauryl Sulfate 2.70%, Inert Ingredients (Water and Sodium Benzoate 96.30%

Key Benefits

Gentle yet effective kills fleas, flea eggs, and ticks on contact, while also repelling mosquitoes.

Certified natural essential oils including peppermint oil and clove extract help control and kill parasitic pests.

Non-staining formula is great for home surfaces including furniture, carpet, flooring, countertops and more.

Plant-based formula contains no harsh chemicals; safe for use around pets, people and even plants.

Proudly made in the USA from high-quality plant-based ingredients and certified natural essential oils.

Precautions

For use with dogs and puppies 12 weeks or older. Keep out of the reach of children and pets.

Antibiotics

Antibiotics

Gentamicin 250

For Animal Use Only

Gentamicin is used in both dogs and cats to treat or prevent bacterial infections including respiratory infections, wound infections, pneumonia, bloodstream infections, bladder infections and infections of the skin and ear.

Dosage and Administration

Always follow the dosage instructions from your veterinarian.

- The usual dose in dogs is 5 to 7 mg per pound (10 to 15 mg/kg) every 24 hours by intravenous, intramuscular or subcutaneous administration.
- The usual dose for cats 3 to 4 mg per pound (5 to 8 mg/kg) once daily by intravenous, intramuscular or subcutaneous administration.
- When applied topically, the dose and frequency depends on the type of condition being treated. Gentamicin is usually applied two or three times daily.
- The duration of administration depends on the condition being treated, response to the medication and the development of any adverse effects. Be certain to complete the prescription unless specifically directed by your veterinarian. Even if your pet feels better, the entire treatment plan should be completed to prevent relapse or prevent the development of resistance.

Contraindications

If hypersensitivity to any of the components occurs, discontinue treatment and institute appropriate therapy.

Missed Dose

If a dose is missed, apply as soon as you remember. If the dose is close to the next dose, skip the application you missed and resume your regular schedule.

AMOXICILLIN TABLET



INGREDIENT:

Each tablet contains:

Amoxicillin 250mg, 500mg

INDICATIONS

Amoxicillin is a broad-spectrum penicillin antibiotic available by prescription used to treat various infections caused by susceptible strains of bacteria. Amoxicillin can be used to treat infections such as ear infections, urinary tract infections, and skin infections. Amoxicillin may also be used for purposes other than those listed

DIRECTION FOR USE

Dogs/Puppies, Cats/Kittens:

Give 5-10 mg/lb of pet's body weight every 12-24 hours. Your veterinarian will determine the actual dose based on the type of infection being treated.

CAUTIONS

Tell your veterinarian if your pet is being given another antibiotic. Also tell your veterinarian if your pet has allergies, or kidney, stomach or intestinal disease. While symptoms may subside early, use all of the prescribed medicine.

STORAGE

Store below 30°C (room temperature). Protect from light and moisture. Close lid tightly after use

CAUTION Keep lid tightly sealed to preserve freshness. Keep out of reach of children.

PACKING 100 tablets per bottle, 6 tablets/blister

Enrofloxacin tablet

INGREDIENT:

Each tablet contains:

Enrofloxacin 68 mg, 136 mg

INDICATIONS

Enrofloxacin is used in the treatment of many types of bacterial infections including urinary tract, skin, prostate, GI tract, liver, and lungs. Enrofloxacin is the generic equivalent to Baytril and comes in a tasty chewable tablet, which most pets love. Enrofloxacin is sold per tablet and requires a prescription from your veterinarian.

DIRECTION FOR USE

For Dogs: Administer orally at a rate to provide 5-20 mg/kg (2.27 to 9.07 mg/lb) of body weight. For Cats: Administer orally at a rate to provide 5 mg/kg (2.27 mg/lb) of body weight.

Give either as a single dose or divided into 2 equal daily doses administered at 12 hour intervals. The dose should be continued for at least 2-3 days beyond cessation of clinical signs, to a maximum of 30 days.

If you suspect that your pet has received an overdose of Enrofloxacin, consult your veterinarian or emergency veterinary clinic immediately.

STORAGE

Store below 30°C (room temperature). Protect from light and moisture. Close lid tightly after use

CAUTION Keep lid tightly sealed to preserve freshness. Keep out of reach of children.

PACKING 100 tablets per bottle, 6 tablets/blister



METOCLOPRAMIDE

ANTI-EMETIC for Dogs & Cats

DESCRIPTION:

Metoclopramide is classified as an anti-emetic or anti-vomiting drug. Metoclopramide is prescribed to treat a wide variety of stomach issues that include vomiting, nausea, acid reflux disease, or food compaction. Metoclopramide blocks chemicals in the brain that cause your pet to vomit while stimulating the contraction of the stomach and intestines to help move food through the digestive tract.

HOW IT WORKS

When motility (movement) in the stomach is reduced, food pools and creates a sensation of nausea and bloating. In some cases, bile refluxes from the intestine back into the stomach, causing irritation and more nausea. Metoclopramide normalizes stomach contractions so that food and bile can pass in the correct direction. Additionally, metoclopramide is able to cross a biochemical blood-brain barrier to control nausea. Metoclopramide helps the vomiting pet by acting directly on the brain to reduce the sensation of nausea.

COMPOSITION

Each tablet contains Metoclopramide 10mg

DOSAGE AND ADMINISTRATION:

The usual dose is 0.1-0.2mg per pound of pet's body weight every 6-8 hours

STORAGE:

Store below 30°C (room temperature). Protect from light and moisture. Close lid tightly after use.

CAUTION:

Metoclopramide is available by prescription only

Give each dose with plenty of water. Give exactly as directed by your veterinarian.

Keep out of the reach of children and pets.

Do not use in animals with known allergies to metoclopramide.

Do not use Preventic Flea Collars on your pet while giving metoclopramide.

Use caution in animals with kidney damage, diabetes, high blood pressure, or recent stomach surgery.

Notify your veterinarian if your pet is taking selegiline, digoxin, cyclosporine, tetracycline, insulin, narcotic pain relievers, anticholinergic medications, or antispasmodic medications.

Possible Side Effects

Allergic reactions and serious side effects are rare, but in the case of an allergic reaction or a serious side effect, seek immediate veterinary attention. Some common signs of allergic reactions and serious side effects are breathing difficulties, facial swelling, hives, jaundice, or yellowing of the skin, or spasms and seizures.

PACKING

1g/Tab x 120 Tab/Bottle x 70 Bottles/Cartron

Analgesics

Analgesics

Carprofen

Carprofen is a non-steroidal anti-inflammatory (NSAID) used to treat pain and inflammation due to arthritis and other joint diseases, including hip dysplasia. It's also used to reduce pain associated with surgery or dental procedures, as well as to reduce fever. Carprofen works by reducing hormones that cause pain and inflammation in the body. Carprofen may also be used for purposes other than those listed here. Carprofen requires a prescription from your veterinarian.

For:

Dogs (over six weeks of age)

Weight:

All weights

DOSAGE AND ADMINISTRATION:

The recommended dosage for oral administration to dogs is 2 mg/lb of body weight daily. The total daily dose may be administered as 2 mg/lb of body weight once daily or divided and administered as 1 mg/lb twice daily. For the control of postoperative pain, administer approximately 2 hours before the procedure.

**Tip:**

Carprofen should be given with food.
Keep plenty of water available for your pet to drink.
Carprofen is for dogs only. Do not give to cats.

Cautions:

Talk to your veterinarian or pharmacist before giving any other prescription or over-the-counter medications to your pet. Should not be given to pregnant or lactating dogs or dogs used for breeding.

Tell your veterinarian if your pet has: an allergy to aspirin or any other NSAID; an ulcer or bleeding in the stomach; liver or kidney disease; congestive heart failure; fluid retention; bleeding disorders; heart disease; high blood pressure.

Nutritional Supplements

Nutritional Supplements



LIVER CARE

Liver Supplement

for DOGS

DESCRIPTION:

LIVER CARE chewables maintain normal liver health and function in dogs. The combination of milk thistle, vitamin E and zinc together with B complex vitamins and taurine in easy-to-give liver flavor chewables provide a multi-faceted approach to liver support. Ideal for older dogs.

COMPOSITION

N-Acetyl L-Cysteine	20 mg
Milk Thistle Extract	10 mg
Vitamin E (dl Tocopheryl Acetate)	10 IU
Zinc (Oxide)	2 mg
L-Taurine	10 mg
Pyridoxine	8 mg
Thiamine (Minonitrate)	8 mg
Riboflavin	4 mg
Niacinamide	8 mg
Vitamin B12	3 mcg



Inactive Ingredients:

Beef Liver, Magnesium Silicate, Magnesium Stearate, Natural Pork Flavor, Plant Cellulose, Pork Liver, Silicon Dioxide, Stearic Acid, Sucralose

DOSAGE AND ADMINISTRATION:

One (1) tablet twice daily per 20 lbs body weight.

PACKING

60 Tablet per bottle or as your requirement

STORAGE:

Store below 30°C (room temperature). Protect from light and moisture. Close lid tightly after use.

CAUTION:

For animal use only. Keep pout of the reach of children and other animals. In case of accidental overdose, contact a veterinarian immediately. Do not use if product appears tampered with or seal is broken.

Bone Live (Glucosamine plus)



DESCRIPTION:

Bone Live will assist the joint mobility of senior animals- dogs & cats. These tablets combines joint repairing supplements- glucosamine and chondroitin - helping to remain or repair the health of your pets joints.

*Chondroitin Sulphate is the major Glycosamine Glycan (GAG) found in the cartilage.

*MSM is an ideal source of bioavailable sulphur.



COMPOSITION

Glucosamine Sulphate.....500mg
Chondroitin Sulphate.....250mg Green Lipped Mussel (stabilized).....100mg
Manganese Ascorbate.....90mg MSM (Methylsulfonyl Methane).....100mg

DOSAGE

Dosage for first 4 Weeks of Use (Dogs & Cats) Up to
5kg.....1/2 Tablet
5kg to 10kg.....1 Tablet
10kg to 20kg.....2 Tablets
20kg to 30kg.....3 Tablets
30kg to 40kg.....4 Tablets

Maintenance Dose

Up to 5kg.....1/4 Tablet
5kg to 10kg.....1/2 Tablet
10kg to 20kg.....1 Tablet
20kg to 30kg.....1 1/2 Tablets
30kg to 40kg.....2 Tablets

DIRECTIONS FOR USE

Give half a dose in the morning and half a dose in the evening. Tablet can be given whole or crushed and mixed with water.

PACKING

60 Tablet per bottle or as your requirement

STORAGE:

Store below 30°C (room temperature). Protect from light and moisture. Close lid tightly after use.

Healthy Coat Omega 3-6



ACTIVE INGREDIENT PER 2g

Crude Protein	35%
Crude Fat	5%
Crude Fiber (max)	5.5%
Moisture (min).....	10%
Calcium (min).....	2%
(Max).....	3%
Phosphorous.....	2%
Linoleic Acid (omega 6)	0.5%
Linoleic Acid(Omega 3)	0.4%

Inactive Ingredients:

Dry whey, Di-calcium phosphate, cellulose, Flaxseed Oil, Sunflower Oil.

Indication:

Healthy Coat Omega 3 & 6 is veterinarian recommended to support skin and coat health in dogs with food or environment sensitivities or seasonal allergies. Our great tasting chewables contains omega 3 and omega 6 fatty acids (EPA, DHA and GLA) which become catalyst for healthy skin and glossy coat in dogs. Works quickly to support a soft, silky coat and minimize normal shedding.

Dosage:

Weight	
10 LBS	1g twice daily
20 LBS	2g twice daily

Net weight: 120g

WARNING: For Animal Use Only.

Keep out of reach of children. Do not leave product unattended around pets. In case of overdose, contact your vet immediately.

Cat Hairball Remedy Gel Supplement

safe, gentle, easy-to-use gel

Hairball Remedy Tuna Flavor Gel Cat Supplement is a smooth easy-to-use gel for the prevention and removal of hairballs in cats and kittens. Both long and short haired cats shed to some degree all year long. Bathing and brushing helps lower the amount of hair swallowed during self-grooming, but some hair gets into the stomach and it cannot be digested. It can block your cat's digestive system. Regular use of Cat Hairball Remedy Gel Supplement the swallowed hair and allows it to easily pass through the intestinal tract.

Key Benefits

- Gentle but effective lubricant for the elimination and prevention of hairballs
- Helps eliminate hairball in the litter box rather than create a mess on the floor
- Safe and effective hairball treatment may be used daily to prevent and treat hairballs
- Regular use coats swallowed hairballs and allows them to pass through the intestinal tract
- Easy to use, palatable gel makes administering to your cat easy and enjoyable

Ingredients

White Petrolatum Light Mineral Oil Nf, Corn Syrup, Malt Syrup, Soybean Oil, Cane Molasses, Water, Gelatin By-Products, Sodium Benzoate (Preservative), and Natural and Artificial Flavors.

For Cat Hairballs: Adult Cats: Administer 1/2 teaspoon daily. Kittens Over 4 Weeks: Administer 1/4 teaspoon daily.

To Prevent Hairballs: Administer 1/2 teaspoon once or twice per week.

Store in a cool, dry place.

Precautions

HAZARDS TO ANIMALS: Safe use in pregnant animals or animals intended for breeding has not been proven. If animal's condition worsens or does not improve, stop product administration and consult your veterinarian. **WARNING:** For animal use only. Keep out of reach of children and animals. In case of accidental overdose, contact a veterinarian immediately.

Shake well. Administer orally. Place a small amount on cat's nose or paws to stimulate taste interest.



Probiotic-vita Paste Daily Supplement For Dog and Puppies



Ingredient

Inactivated Lactobacillus bulgaricus, Fructooligosaccharide, Vitamin A, Vitamin B1, Vitamin B2, Vitamin B6, Vitamin B12, Vitamin D, Vitamin E, Calcium Phosphate, Ferrous Sulfate, Zinc Sulfate, Chicory root powder, oligo-maltose, Leguminous seed

Each 100g contain

Vitamin A.....	9600IU
Vitamin D.....	600IU
Vitamin E.....	90 mg
Vitamin B1.....	6.5 mg
Vitamin B2.....	7.0 mg
Vitamin B6.....	3.8 mg
Vitamin B12.....	30 ug
Nicotinic acid.....	20 mg
Choline.....	300 mg
Folic acid.....	200 ug
Pantothenic acid.....	12 mg
Biotin.....	160 ug
Calcium.....	0.2 g
Phosphorus.....	0.18 g
Iron.....	3.6 mg
Zinc.....	3.7 mg
Inactivated Lactobacillus bulgaricus.....	1 x 10 ⁹ cfu

Benefits

Coordinating intestines and stomach, fast recovering the digestive system, maintaining the health of intestines and stomach.

Promote appetite, restore physical strength.

Help to eliminate halitosis caused by gastrointestinal disorders.

Comprehensive nutrition supplies, rapidly improve the intestinal power.

Recommended Dosage

Feed directly or add food

5kg ~ 10kg 5 ~ 8cm/day

10kg ~ 25kg 8 ~ 10cm/day

Over 25kg 10 ~ 12cm/day

Net Weight: 120g

Caution: Keep Out of reach of children.

Keep container tightly sealed, store in a cool dry place below 25 C



Chewable Calcium Calcium Supplement for Dog&Cat

VIC CAL

Chewable Calcium, an excellent growth promoter in Dogs & Cats for all age groups fortified with calcium, phosphorus, vitamin D3, biotin and essential amino acids.

COMPOSITION

Calcium	83 mg
Phosphorus	34 mg
Vitamin D3	200 IU
Vitamin B12	5 mcg
Biotin	5 mcg
Magnesium	0.32 mg
Lysine	4 mg
Methionine	2 mg
Protein	48 mg

INDICATIONS:

CHEWABLE CALCIUM has been specially formulated as a source of calcium in young as well as geriatric pets. The unique composition of the supplement helps prevent rickets, osteoporosis, osteomalacia in pets. It also aids in faster recovery and healing of fractures and promotes healthy bones and good growth.

DOSAGE AND ADMINISTRATION:

Small Dog / Cats - 1 tab twice a day Medium Dog / Cats - 2 tabs twice a day Large and giant breeds - 4 tabs twice a day

STORAGE:

Store below 30°C (room temperature). Protect from light and moisture. Close lid tightly after use.

CAUTION:

For animal use only. Keep out of the reach of children and other animals. In case of accidental overdose, contact a veterinarian immediately. Do not use if product appears tampered with or seal is broken.





Chewable Vitamin Liver Supplement for DOGS

Nutritional Supplements



VIC CAL PLUS

Chewable Vitamin is more than an tasty all-natural multi-vitamin, it is the ultimate blend of acids, vitamins and minerals. Together these natural ingredients support a healthy immune system and circulatory functions thereby promoting optimal health in your pet.

Chewable Vitamin contains 21 amino acids, 11 minerals and 14 vitamins. The addition of these 21 amino acids makes Chewable Vitamin truly unique and revolutionary in the world of multi-vitamins for pets.

INGREDIENTS:

Dextrates, Maltodextrin, Whey, Natural Flavorings, Stearic Acid, Dicalcium Phosphate, Silicon Dioxide, Potassium Chloride, Cellulose, Gum Ghatti, Egg Albumin, L-Carnitine, L-Taurine, Sunflower Oil, Ferrous Fumarate, Choline Bitartrate, DL-alpha-Tocopheryl Acetate, Fish Protein Concentrate, Ascorbic Acid, Zinc Acetate, Magnesium Oxide, Inositol, Pantothenic Acid, Niacin, Vitamin A Palmitate, Riboflavin, Thiamine Mononitrate, Pyridoxine Hydrochloride, Manganese Gluconate, Vitamin D3, Copper Gluconate, Potassium Iodide, Sodium Selenate, Vitamin B12, Biotin, Folic Acid.

Active Ingredients per 2g:

INDICATIONS:

Improving health and vitality

DOSAGE AND ADMINISTRATION:

Give as treat or crumble and mix with food according to the following schedule:

Small Dogs (less than 20 lbs.): 1 tablet daily. Mid-sized Dogs (20-40 lbs.): 2 tablets daily. Larger Dogs (41-60 lbs.): 3 tablets daily.

Large Dogs (61-80 lbs.): 4 tablets daily

Very Large Dogs (81-100 lbs.): 5 tablets daily. Giant Breeds (100-150 lbs.): 6-7 tablets daily.

STORAGE:

Store below 30°C (room temperature). Protect from light and moisture. Close lid tightly after use

PACKING

120g per bottle

CAUTION:

For animal use only. Keep out of the reach of children and other animals.

Amino Acids	
Arginine	19.9 mg
Histidine	14.4 mg
Isoleucine	43.7 mg
Leucine	74.9 mg
Phenylalanine	22.8 mg
Tyrosine	21.2 mg
Methionine	15.0 mg
Cystine	18.7 mg
Threonine	49.2 mg
Tryptophan	12.1 mg
Valine	40.4 mg
Lysine	65.5 mg
Alanine	29.6 mg
Aspartic acid	75.8 mg
Glutamic acid	118 mg
Glycine	13.9 mg
Proline	44.2 mg
Serine	38.4 mg
L-Carnitine	15.0 mg
Taurine	15.0 mg
Panthenic acid	540 mcg
Niacinamide	540 mcg
Folic acid	3.6 mcg
Choline	2.16 mg
Inositol	1.08 mg
Biotin	4.5 mcg

Minerals	
Calcium	21.6 mg
Phosphorus	18.0 mg
Iron	1.8 mg
Copper	54 mcg
Zinc	1.35 mg
Iodine	41.4 mcg
Potassium	28.6 mg
Manganese	54 mcg
Selenium	2.97 mcg
Cobalt	1.8 mcg
Magnesium	900 mcg

Vitamins	
Vitamin A	144 IU
Vitamin D3	10.8 IU
Vitamin E	1.44 IU
Vitamin C	1.8 mg
Vitamin B1	180 mcg
Vitamin B2	226 mcg
Vitamin B6	180 mcg
Vitamin B12	9 mcg

Pet Milk Supplements



Nutritional Supplements

Main effects:

The formula uses the highest quality natural milk source, goat milk, which is closest to breast milk, to provide rich and low-sensitive nutrient molecules;

Add whey protein to improve the immune resistance of puppies/cats and quickly supplement the nutrients required for dogs/cats at each stage of growth.

Probiotics + prebiotics regulate the intestinal micro-ecological balance, promote the digestion and absorption of nutrients in the intestines, and provide special love and care for pets.

Hypoallergenic factor: Casein is a common allergic factor in dairy products. The casein content in goat milk is 90% less than that of milk, which can reduce the allergic reaction caused by ingestion of dairy products in dogs/cats.

Low-lactose formula: goat milk contains low lactose content, and lactase is added to prevent soft stools and diarrhea in dogs/cats due to lactose intolerance.

Main ingredients:

Raw material composition: goat milk powder, whole milk powder, whey protein, maltodextrin, glucose

Additive composition: oligosaccharide, Bifidobacterium lactis, milk calcium, multivitamin (vitamin A, vitamin D3, vitamin E, vitamin B1, vitamin B2, vitamin B6, vitamin B12, niacin, folic acid, pantothenic acid, choline chloride), complex trace elements (iron, zinc, copper, iodine, calcium, phosphorus), lactase

Guaranteed nutrition:

Nutrition ingredient	Content	Nutrition ingredient	Content	
Energy (ME)	KJ/100g	1900	Niacin acid mg	≥ 7.50
Crude protein %	≥	24.0	Pantothenic acid mg	≥ 6.00
Crude fat %	≥	15.0	Vitamin B6 mg	≥ 2.00
Crude fibrin %	≤	10.0	Folic acid mg	≥ 0.10
moisture %	≤	6.0	Vitamin B12 mg	≥ 0.015
Crude ash %	≤	8.0	Choline mg	≥ 140
Calcium %	≥	2.40	Potassium %	≤ 0.06
Lysine (dog) %	≥	0.63	Chlorine %	≥ 0.10
Taurine (cat) %	≥	0.1	Magnesium %	≥ 0.024
Probiotics cfu	≥	1.0×10 ¹¹	Ferrite mg	≥ 10
Vitamin A IU	≥	5000	Copper mg	≥ 1.00
Vitamin D3 IU	≥	500	Manganese mg	≥ 1.00
Vitamin E mg	≥	15	Zinc mg	≥ 12.00
Vitamin B1 mg	≥	2.50	iodine mg	≥ 0.15
Vitamin B2 mg	≥	3.00	Selenium mg	≥ 0.020

Applicable object/product function:

Newborn puppies: balanced and complete nutrition, can replace breast milk, enhance disease immunity, promote growth and development.

Pregnant lactating bitch cat: supplement nutrition, promote healthy embryo development and increase milk production.

Anorexia / Post-healing rehabilitation / old and weak dogs and cats: rapid nutritional supplements, promote appetite and enhance physical strength.

Supplement or replace breast milk stage

Body Weight	Recommended feed	Feeding frequency
≤120g	60ml/day	0~2 weeks old: feeding every 2 hours
120g~160g	70~100ml/day	
160~210g	100~130ml/day	2~4 weeks old: feeding every 3 hours
210g~280g	130~170ml/day	Full moon or more: feeding every 4 hours
≥280g	200ml/day	

Supplementary stage

Stage	Recommended feed
Weak anorexia	100ml/day
Gestation period	100ml/once, twice per day
Lactation period	
Postmastectomy relief period	100ml/once, thrice per day
Gerontic(Over 6 age)	

Usage: Every 10 grams of milk powder plus 60 °C warm water 60ml brewing to be cold to 37 °C or so to feed directly.

Shelf life: 24 months

Storage conditions: Please store in a cool and dry place below 25 °C., avoid direct sunlight, please eat as soon as possible after opening.

Note: This product is limited to animals, please go to the location of the product that children can not touch.

Digestive Support

COMPOSITION

Active Ingredients: Amounts Per Scoop (2.5 Grams)

Proprietary Digestive Blend 113.95mg

Enzymes: Protease, Lipase, Alpha-galactosidase, Amylase, Phytase, Lactase, Invertase and Hemicellulase

Probiotics: Lactobacillus acidophilus, Bifidobacterium bifidum, Lactobacillus casei, Lactobacillus plantarum & Lactobacillus reuteri

Digestive Support Ingredients:

Fructooligosaccharides 145.61mg Ginger Root (Zingiber officinale) 48.54mg

Lemon balm (Melissa officinalis) 50.48mg Psyllium Husk (Plantago ovata) 9.71mg

Inactive Ingredients: Beef protein Isolate, chicken liver, chicken protein isolate, whey protein Isolate

DESCRIPTION

Digestive Support is a highly potent supplement powder that provides a concentrated level of essential nutrients for the support and maintenance of a healthy digestive system. It contains a unique enzyme blend, powerful ingredients known for absorbing and removing harmful intestinal toxins, and a complex system of pre- and probiotics for dogs.

Mainly Support For: Nutrient Digestion, Nutrient Absorption, Optimum Nutrient Utilization, Microflora Health, Nutrient Digestion, Nutrient Absorption, Optimum Nutrient Utilization, Microflora Health

PRECAUTIONS

Use with caution in pregnant or lactating dogs. Do not use in dogs that have been diagnosed with diabetes, gallstones or intestinal blockage unless advised by a veterinarian.

DOSAGE AND ADMINISTRATION

Recommend for growing, adult, and senior dogs.

Sprinkle dry onto any canine diet. This product may also be moistened to make a tasty broth.

Dogs (All ages)	Weight Range	Scoops/Day (2.5g per scoop)
Toy	3 - 12 LBS	1/3
Small	13 - 25 LBS	2/3
medium	26 - 44 LBS	1
Med-large	45 - 69 LBS	1 (1/2)
LARGE	70 - 99 LBS	2
GIANT	Over 100 LBS	2 (1/2)

As with any change in your dog's diet, it's very important to start slowly. Begin by giving your dog one tablet daily with meals for at least 2 - 3 days. This will allow your pet to adjust gradually to the change.

STORAGE

Store in a cool, dry area out of direct sunlight. Keep lid tightly sealed to preserve freshness.

Turmeric chewable tablet

COMPOSITION

Each 1.0g tablet contains.

Turmeric Root Extract (Curcuma longa L) 300mg

Other Ingredients: Microcrystalline Cellulose, Dicalcium Phosphate, Liver (Dessicated Beef, Undefatted), Stearic Acid, Bacon Flavor, Croscarmellose Sodium, Magnesium Stearate, Silicon Dioxide.

DESCRIPTION

The Turmeric Chews Daily Dog Supplement has curcumin, a proven and powerful antioxidant naturally derived from 100% natural turmeric root to help soothe the painful joint inflammation and promote a healthy heart, liver and digestion. The hypoallergenic and veterinarian-recommended formula is easy to give with no messes, hassles, or smells. Each chewable tablet contains pharmaceutical grade curcumin, and is naturally flavored with bacon and liver so your pal will love them as a treat! They're ideal for dogs with hip and joint discomfort from arthritis or everyday wear and tear.

KEY BENEFITS

Helps soothe painful joint inflammation and promotes liver, heart, and digestive health.

Formulated with curcumin, a powerful antioxidant naturally derived from 100% natural turmeric root.

Hypoallergenic formula is veterinarian-recommended for dogs with joint discomfort from arthritis or everyday activity.

Naturally flavored with bacon and liver and deliciously chewable so your dog will love them as a daily treat. No mess or smells, no wheat, corn, maltodextrin, animal digest, grains, soy, egg, potato, or known allergens.

PRECAUTIONS

If too much is taken, side effects are generally minimal and may include diarrhea. Always consult your veterinarian. Keep out of reach of children.

DOSAGE AND ADMINISTRATION

Recommended Usage: 2 tablets daily, depending on your dog's weight and individual needs. Allow 3 - 4 weeks to note a response, some dogs may respond sooner. Best fed with food.

Weight	Daily Guide
UNDER 40 LBS	2 tablet per day (can be split in half for smaller dogs)
41 - 80 LBS	3 - 4 tablets per day
81 - 120 LBS	5 - 6 tablets per day

As with any change in your dog's diet, it's very important to start slowly. Begin by giving your dog one tablet daily with meals for at least 2 - 3 days. This will allow your pet to adjust gradually to the change.

STORAGE

Store in a cool, dry area out of direct sunlight. Keep lid tightly sealed to preserve freshness.

Pet Blood producer nutritional supplements



Ingredient: hemoglobin powder, chicken liver powder, astragalus, angelica, brewer's yeast powder, fish oil.

Additive composition: ferrous gluconate, taurine, lecithin, vitamin A, vitamin B1, vitamin B2, vitamin B6, vitamin B12, vitamin C, vitamin D3, vitamin E, zinc sulfate, magnesium sulfate

Analysis guaranteed value

Ingredient	Per kg contain
crude protein	≥16%
crude fat	≥15%
moisture	≤10%
crude ash	≤5%
crude fibre	≤2%
taurine	2500mg/kg
vitamin A	2800IU/kg
vitamin B6	10mg/kg
vitamin B12	0.1mg/kg
Folic acid	0.6mg/kg
vitamin D3	1000IU/kg
vitamin E	200mg/kg
calcium	0.1%
phosphorus	0.08%
iron	377mg/kg
Zinc	16.5mg/kg
magnesium	18mg/kg



Product Description:

Heme protein powder is rich in protein and heme iron. Heme iron can be directly absorbed in intestinal mucosal epithelial cells, and the iron absorption and absorption rate is high. Angelica and Astragalus polysaccharide extract, adjust the vitality and nourish the blood. B vitamins It is a component of coenzyme, which helps to improve hematopoietic function, enhance cell activity and function, increase appetite, and help the body to recover. It also adds taurine, multivitamins and trace elements to supplement nutrition and enhance the health of pets. benefiting energy, supplement iron and Produce blood effect; suitable for iron deficiency anemia, excessive blood loss, nutritional imbalance caused by anemia.

Dosage and usage:

Applicable to dogs/cats with iron deficiency anemia, excessive blood loss, and nutritional imbalance caused by anemia. This product is palatable, can be fed directly added to food or crush.

Puppies and cats ≤5kg 2 capsules / day

Small dog 5-10kg 3-4 capsules / day

Medium dog 10-25kg 4-6 capsules / day

Large dogs 25-40kg or more 6-8 tablets / day

Note: This product should not be fed ruminants, please keep out of reach of children.

Storage method: Please store in a cool and dry place below 25 °, and avoid direct exposure to sunlight.

Shelf life: 24 months



Terbinafine Spray

Terbinafine is a medication used to treat fungal infections especially those of the skin, hair, and nails in cats and dogs. It works by stopping the growth of fungal cells and killing those fungal cells.

How Terbinafine Works

Terbinafine is classified as an allylamine antifungal antibiotic. Allylamine antifungal antibiotics work by inhibiting the synthesis of an enzyme called squalene epoxidase in fungi. Fungi use this enzyme to synthesize another chemical, called ergosterol, which is used in the cell membrane of fungal cells. By keeping the fungus from producing squalene epoxidase, allylamine antifungal antibiotics indirectly prevent the fungus from building and maintaining cell membranes. Eventually, this will weaken the fungal cells enough to cause ruptures to appear in their cell membranes (a process called lysis), which allows the cell's contents to spill out, thus killing the cell. Once terbinafin enters a cat's body, it tends to accumulate in the skin, hair, claws and fatty tissue. This tendency of Terbinafine to accumulate around the skin and hair makes it an ideal treatment for ringworm infections.

Ringworm in Cats

Ringworm is caused by a fungal infection in the skin that results in hairless, circular lesions that are often covered in pustules. These lesions are usually very irritated and itchy so they may be red and inflamed, possibly with further damage caused by excessive scratching. Ringworm lesions in cats are most likely to appear on the head, ears, and the base of the tail.

Side Effects

Terbinafine may cause some side effects in dogs. Because it hasn't been properly tested, it's a good idea to keep careful track of these effects and to notify your vet immediately. The side effects include:

- Vomiting and diarrhea
- Swelling of the abdomen
- Faintness or dizziness
- Nausea
- Loss of appetite
- Weight loss

side effect in cats: vomiting.

If you recognize any of these side effects, it could be the symptom of either an allergic reaction or some other issue that is a result of your pet taking the terbinafine supplement and medicine. Take your dog to the vet right away at the first sight of these effects.

Terbinafine can be a great way of helping to deal with your pet's external fungal infections. Left untreated, these can spread to other parts of your dog's body and cause him not only great discomfort, but also more serious health concerns. Eliminate the fungus fast with terbinafine treatment.

Dosage: 30-40mg/kg q24 hours

HEARTWORM REMEDY PLUS

Ivermectin Plus Pyrantel Pamoate Tablet

COMPOSITION

[FOR SMALL SIZE DOGS] Each Chewable Tablet Contains:

Ivermectin BP 68 mcg
 Pyrantel Pamoate USP equivalent to Pyrantel 57 mg

[FOR MEDIUM SIZE DOGS] Each Chewable Tablet Contains:

Ivermectin BP 136 mcg
 Pyrantel Pamoate USP equivalent to Pyrantel 114 mg

Excipients q.s.

[FOR LARGE SIZE DOGS] Each Chewable Tablet Contains:

Ivermectin BP 272 mcg
 Pyrantel Pamoate USP equivalent to Pyrantel 227 mg

INDICATIONS

For use in dogs to prevent canine heartworm disease by eliminating the tissue stage of heartworm larvae (*Dirofilaria immitis*) for a month (30 days) after infection and for the treatment and control of ascarids (*Toxocara canis*, *Toxascaris leonina*) and hookworms (*Ancylostoma caninum*, *Undnaria stenocephala*, *Ancylostoma braziliense*).

DOSAGE

Orally at monthly intervals at the recommended minimum dose level of 6 mcg of Ivermectin per kilogram (2.72 mcg/lb) and 5 mg of Pyrantel (as pamoate salt) per kg (2.27 mg/lb) of body weight. The recommended dosing schedule for prevention of canine heartworm disease and for the treatment and control of ascarids and hookworms is as follows:

Dog Weight		Tablet Per Month	Ivermectin Content	Pyrantel Content
kg	lbs			
Upto 11kg	Upto 25 lbs	1	68 mcg	57 mg
12-22kg	26-50 lbs	1	136 mcg	114 mg
23-45kg	51-100 lbs	1	272 mcg	227 mg

This product is recommended for dogs 6 weeks of age and older.

For dogs over 100 lbs use the appropriate combination of these Chewable Tablets.

ADMINISTRATION

This product should be given at monthly intervals during the period of the year when mosquitoes (vectors), potentially carrying infective heartworm larvae, are active. The initial dose must be given within a month (30 days) after the dog's first exposure to mosquitoes. The final dose must be given within a month (30 days) after the dog's last exposure to mosquitoes.

EFFICACY

Heartworm Remedy Tablet, given orally using the recommended dose and regimen, are effective against the tissue larval stage of *D. immitis* for a month (30 days) after infection and, as a result, prevent the development of the adult stage. Strongheart Plus chewable tablets are also effective against canine ascarids (*T. canis*, *T. leonina*) and hookworms (*A. caninum*, *U. stenocephala*, *A. braziliense*).

ACCEPTABILITY

In acceptability and field trials, Strongheart Plus was shown to be an acceptable oral dosage form that was consumed at first offering by the majority of dogs.

STORAGE:

Store at controlled room temperature (15-30°C). Protect from light.

PACKING

10 Tablets/ Blister, 10 Blisters/Box, 40 Boxes/ Carton
 100 Tablets/Satchel, 1 Satchel/Jar, 60 Jars/ Carton

Manufacturer: Hebei Weierli Animal Pharmaceutical Co Ltd
Phone: +86-311-89265710
Fax: +86-311-68052865
Address: Luquan District, Shijiazhuang City, Hebei Province
Website: <http://www.victorypharm.com/>